


## DOSING GUIDE

This guide refers to dosing of patients prescribed TECENTRIQ as per the EU Summary of Product Characteristics (SmPC). Treatment with TECENTRIQ must be initiated and supervised by physicians experienced in the treatment of cancer. For TECENTRIQ approved indications and full administration support see the SmPC.

### STANDARD TECENTRIQ ADMINISTRATION

- TECENTRIQ is for IV use (**not** IV push or bolus).
- Initial TECENTRIQ dose must be administered over 60 mins. If well tolerated, subsequent infusions may be administered over 30 mins.
- Dose reductions of TECENTRIQ are not recommended.
- If a planned dose of TECENTRIQ is missed, it should be administered as soon as possible.
- TECENTRIQ treatment is recommended until loss of clinical benefit or unmanageable toxicity.
- For pre-treatment care, pre-medication, lab tests, recording of vital signs and follow up, refer to local guidelines and individual SmPCs.

### IMMUNE-RELATED ADVERSE REACTIONS



### INFUSION-RELATED REACTIONS

The rate of infusion should be reduced or treatment should be interrupted in patients with Grade 1 or 2 infusion-related reactions. TECENTRIQ should be permanently discontinued in patients with Grade 3 or 4 infusion-related reactions.

Published by F. Hoffmann-La Roche Ltd, 4070 Basel, Switzerland | © December 2020 www.roche.com

All trademarks mentioned herein are protected by law. M-XX-00003869

<b>TNBC 1L</b>  <b>TECENTRIQ + NAB-PACLITAXEL</b> Treatment recommended until disease progression or unmanageable toxicity.	Three flexible & convenient fixed dosing options. 4-week interval 1680 mg IV (2x 840 mg IV) 3-week interval 1200 mg IV 2-week interval 840 mg IV ■ TECENTRIQ	<b>TECENTRIQ</b>  30 minutes 60 min for 1 <sup>st</sup> infusion	<b>DAYS 1, 8, &amp; 15</b> <b>NAB-PACLITAXEL</b> 100 mg/m <sup>2</sup>  30 minutes	<b>SCLC</b>  <b>TECENTRIQ + CARBOPLATIN + ETOPOSIDE</b> Treatment recommended until disease progression or unmanageable toxicity.	Three flexible & convenient fixed dosing options. 4-week interval 1680 mg IV (2x 840 mg IV) 3-week interval 1200 mg IV 2-week interval 840 mg IV ■ TECENTRIQ	<b>INDUCTION   Q3W</b> <b>TECENTRIQ</b>  30 minutes 60 min for 1 <sup>st</sup> infusion	<b>CARBOPLATIN AUC</b> 5 mg/ml/min <sup>a</sup>  30-60 minutes <sup>a</sup> Dose may vary as per local protocol and/or renal function	<b>MAINTENANCE   Q2W, Q3W or Q4W</b> <b>ETOPOSIDE</b> 100 mg/m <sup>2</sup>  60 minutes same dose administered on Days 2 and 3 as well	<b>TECENTRIQ</b>  30 minutes				
	<b>NSCLC 1L</b>  <b>TECENTRIQ + AVASTIN + PACLITAXEL + CARBOPLATIN</b> Treatment recommended until disease progression or unmanageable toxicity.	Three flexible & convenient fixed dosing options. 4-week interval 1680 mg IV (2x 840 mg IV) 3-week interval 1200 mg IV 2-week interval 840 mg IV ■ TECENTRIQ	<b>INDUCTION   Q3W</b> <b>TECENTRIQ</b>  30 minutes 60 min for 1 <sup>st</sup> infusion		<b>AVASTIN</b> 15 mg/kg  30 minutes	<b>PACLITAXEL</b> 200 mg/m <sup>2</sup>  180 minutes	<b>CARBOPLATIN AUC</b> 6 mg/ml/min <sup>a</sup>  15 - 30 minutes <sup>a</sup> Dose may vary as per local protocol and/or renal function	<b>MAINTENANCE   Q3W</b> <b>TECENTRIQ</b>  30 minutes	<b>AVASTIN</b> 15 mg/kg  30 minutes	<b>NSCLC 1L</b>  <b>TECENTRIQ + NAB-PACLITAXEL + CARBOPLATIN</b> Treatment recommended until disease progression or unmanageable toxicity.	Three flexible & convenient fixed dosing options. 4-week interval 1680 mg IV (2x 840 mg IV) 3-week interval 1200 mg IV 2-week interval 840 mg IV ■ TECENTRIQ	<b>INDUCTION   Q3W</b> Day 1 <b>TECENTRIQ</b>  30 minutes 60 min for 1 <sup>st</sup> infusion	<b>DAYS 1, 8 &amp; 15</b> <b>NAB-PACLITAXEL</b> 100 mg/m <sup>2</sup>  30 minutes
<b>NSCLC 2L</b>  <b>TECENTRIQ</b>	Three flexible & convenient fixed dosing options. 4-week interval 1680 mg IV (2x 840 mg IV) 3-week interval 1200 mg IV 2-week interval 840 mg IV ■ TECENTRIQ	<b>DOSING OPTIONS   Q2W, Q3W or Q4W</b> <b>TECENTRIQ</b>  30 minutes 60 min for 1 <sup>st</sup> infusion Until loss of clinical benefit or unmanageable toxicity.	<b>UC</b>  <b>TECENTRIQ</b>	Three flexible & convenient fixed dosing options. 4-week interval 1680 mg IV (2x 840 mg IV) 3-week interval 1200 mg IV 2-week interval 840 mg IV ■ TECENTRIQ	<b>DOSING OPTIONS   Q2W, Q3W or Q4W</b> <b>TECENTRIQ</b>  30 minutes 60 min for 1 <sup>st</sup> infusion Until loss of clinical benefit or unmanageable toxicity.	<b>HCC</b>  <b>TECENTRIQ + AVASTIN</b> Q3W is the recommended dosing frequency for TECENTRIQ combination therapies.	4-week interval 1680 mg IV (2x 840 mg IV) 3-week interval 1200 mg IV 2-week interval 840 mg IV ■ TECENTRIQ	<b>DOSING OPTIONS   Q3W</b> <b>TECENTRIQ</b>  30 minutes 60 min for 1 <sup>st</sup> infusion	<b>AVASTIN</b> 15 mg/kg  30 minutes 90 min for 1 <sup>st</sup> infusion 60 min for 2 <sup>nd</sup> infusion				

This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions.

<b>TECENTRIQ SC approved in the EU for all indications of intravenous TECENTRIQ<sup>1*</sup></b> 	<b>SCHEDULING</b> Every 3 weeks 	<b>PREPARATION</b> Fixed dose 1875 mg SC (1 x 1875 mg vial) 	<b>ADMINISTRATION</b> Approximately 7 minutes for the first and subsequent injections 
	<p>* including certain types of lung, liver, bladder and breast cancer. Please refer to the TECENTRIQ SmPC for further information</p>		